



# COMMUNITY AIR MONITORING REPORT (FEBRUARY 2019-MARCH 2020)

**Community of South Central Fresno**

San Joaquin Valley Air Pollution Control District

May 27, 2020

# TABLE OF CONTENTS

I.	Community Air Monitoring and Status of Network Deployment .....	1
II.	Summary of PM2.5 Air Monitoring .....	7
III.	Summary of Data Collected using Mobile Air Monitoring Van .....	10
IV.	Summary of PM2.5 Speciation Analysis .....	12
V.	Summary of VOC Speciation Analysis.....	14
VI.	Availability and Access to Community Air Monitoring Data .....	16
VII.	Next Steps .....	16
VIII.	Appendix A: Daily PM2.5 Averages .....	17
IX.	Appendix B: Summary of Data Collected using Mobile Air Monitoring Van .....	28


---

## I. Community Air Monitoring and Status of Network Deployment

Assembly Bill (AB) 617, signed into law in July 2017, has resulted in a statewide effort to reduce air pollution and improve public health in communities that experience disproportionate burdens from exposure to air pollutants statewide through new community-focused and community-driven actions. AB 617 provides mechanisms and resources to implement community-specific air quality monitoring networks, develop and implement emission reduction programs; improve availability of data and other technical information; and invest substantial funding in the community through voluntary incentive funding measures. South Central Fresno, a densely populated community within the city of Fresno, was selected as a first year community by CARB in September of 2018.

District staff provided assistance to the Community Steering Committee (CSC) members by helping them to develop their recommended air monitoring priorities. The District worked with CSC members as they reviewed and evaluated a variety of different resources, including maps of stationary sources, area sources, mobile sources, prevailing wind direction data, and sensitive receptor locations relative to sources of air pollution within the community. The CSC adopted their official recommendation on June 12, 2019, including the deployment of various air monitoring platforms within the community as identified in Figure 1 below.

**Figure 1 South Central Fresno Community Air Monitoring Network Design**



## **Community Air Monitoring Platforms and Equipment**

The District has been working to site and deploy the following high-precision regulatory grade air monitoring systems and platforms, providing flexible options to meet air monitoring needs for the community air monitoring network:

- **Stand-Alone PM2.5 Monitors:** The District will operate fixed air monitoring analyzers to measure ambient PM2.5. These will be placed in their respective locations for sufficient lengths of time to capture annual and peak PM2.5 pollution trends throughout the community, unless monitoring priorities change and monitor relocation is necessary.



- **Compact Multi-Pollutant Air Monitoring System:** These compact air monitoring systems will operate as semi-mobile platforms. Each platform will be equipped with advanced air monitoring analyzers measuring various pollutants, with the ability to communicate the community-level air quality in real time.



- **Air Monitoring Trailer:** These air monitoring trailer systems will operate as semi-mobile platforms. This platform will be equipped with advanced air monitoring analyzers with the ability to communicate the community-level air quality in real time.



- **Mobile Air Monitoring Van:** The van is ideal for focusing on unmonitored areas of concern and regularly surveying the entire community within short timeframes, allowing for a better understanding of the spatial differences in air quality across the community. The air monitoring van can also be used for measuring pollution from on-road sources, and identifying sources of community-level air pollution. Additionally, the van can be parked in one location for longer periods of time to capture daily or weekly pollution from unmonitored areas within the community.



The air monitoring van is a useful tool for evaluation of a large geographic region, but these platforms are best designed for taking a short-term look at the measured pollutants when and where the monitoring occurred. The fixed and semi-mobile platforms are outfitted with instrumentation that is capable of more accurately measuring daily and long-term variations in pollutant concentrations. The use of both mobile and semi-mobile monitoring platforms will be necessary to capture the full picture of the community's air pollution profile.

These air monitoring systems will provide real-time hourly average readings of the following pollutants:

- **PM<sub>2.5</sub>:** PM<sub>2.5</sub> can be composed of any material that has a diameter of 2.5 microns or less. PM<sub>2.5</sub> can be emitted directly as primary PM<sub>2.5</sub> from various sources or formed secondarily through chemical reactions in the atmosphere.

- **Black Carbon (BC):** Black carbon is also known as soot or elemental carbon and formed during incomplete combustion in fuels, including mobile exhaust (mainly diesel), and wood burning
- **Nitrogen Oxides (NO, NO<sub>2</sub>, NO<sub>x</sub>):** Nitrogen oxides (NO<sub>x</sub>) are chemical compounds formed by the combination of nitrogen and oxygen, and are primarily emitted through the combustion of fossil fuels from mobile and stationary sources
- **Volatile Organic Compounds (VOC):** VOCs are released through the burning of various fuels such as gasoline, wood, coal, or natural gas, and can also be released through the use of solvent based consumer products
- **Ozone:** Ozone is not emitted directly into the air, but is created by chemical reactions between NO<sub>x</sub> and VOC in the presence of heat and sunlight
- **Carbon Monoxide (CO):** CO is a colorless, odorless gas that can be harmful when inhaled in large amounts. The greatest sources of CO to outdoor air are cars, trucks and other vehicles or machinery that burn fossil fuels.
- **BTEX:** BTEX is a specified subset of VOCs containing benzene, toluene, ethylbenzene, and xylene. These chemicals appear naturally in crude oil and can be associated with emissions from petroleum refineries, and petroleum storage and fueling stations.
- **Sulfur Dioxide (SO<sub>2</sub>):** SO<sub>2</sub> is a colorless gas with a pungent odor. Sulfur dioxide is produced largely by fossil fuel combustion.
- **Hydrogen Sulfide (H<sub>2</sub>S):** H<sub>2</sub>S is a colorless gas characterized by its foul odor of rotten eggs and can be smelled at low concentrations. Hydrogen sulfide is often produced from the breakdown of organic matter in the absence of oxygen gas, such as in swamps, sewers, and in the crude oil extraction/refining process.

South Central Fresno community air monitoring also includes the capturing of air samples using canisters and filters that are sent to third party laboratories to be analyzed for VOC and PM<sub>2.5</sub> compounds and species present in the local air.

### ***Status of South Central Fresno Community Air Monitoring Network***

Consistent with the community recommended air monitoring network design, the District is in various stages of installing air monitoring systems in the locations identified in the community air monitoring plan. The progress in implementing the community air monitoring network in South Central Fresno is listed below:

*Roosevelt High School (PM2.5)*

The District has placed a real-time PM2.5 monitor (Met One BAM-1022) on the roof at Roosevelt High School on the corner of Tulare and Barton Avenues. Operation of this analyzer began in March 2019. Data collected from this site is available on the District's South Central Fresno AB 617 air monitoring webpage and is being uploaded to CARB's [AQview portal](#) on a regular basis.

*Bitwise South Stadium (PM2.5)*

The District has placed a real-time PM2.5 monitor (Met One BAM-1022) on the roof at Bitwise South Stadium on the corner of Van Ness Avenue and Mono Street in downtown Fresno. Operation of this analyzer began in August 2019. Data collected from this site is available on the District's South Central Fresno AB 617 air monitoring webpage and is being uploaded to CARB's [AQview portal](#) on a regular basis.

*Fresno-Foundry Park (PM2.5, VOC/PM2.5 speciation)*

The District has placed a real-time PM2.5 monitor (Met One BAM-1020) at the existing District air monitoring site at Foundry Park Ave near the intersection of Jensen Avenue and Highway 99. The PM2.5 analyzer began its official operation in January 2020. The District also began operating VOC and PM2.5 speciation sampling at this location to begin to build an understanding of the relative comparison between the constituents that comprise the VOC and PM2.5 concentrations present in the community. These speciation measurements began in December of 2019. VOC and PM2.5 speciation air monitoring efforts will shift to the trailer at Malaga Elementary School once this site is established.

*Orange Center School (Compact Multi-Pollutant System)*

The District has been in discussions with the Orange Center School District to place an Air Pointer compact multi-pollutant air monitoring system at Orange Center Elementary School on Cherry Avenue. On September 2019, District staff presented a proposal to the Superintendent and School Board. On February 13, 2020, the Superintendent informed District staff that the School Board had voted and denied the request due to a high level of ongoing projects already in progress at the school. District staff will continue to work with the steering committee on this location, and are also working on finding alternative locations and in the interim is using the air monitoring van to monitor areas nearby this location.

*Heaton Elementary School (PM2.5)*

The District has been in discussion with Fresno Unified School District (FUSD) in regards to placing a real-time PM2.5 monitor at Heaton Elementary School on the corner of San Pablo and McKinley Avenues. The District is continuing to attempt to schedule further meetings with FUSD to discuss implementation details. In the interim,

the air monitoring van is being utilized to monitor areas nearby the school. In addition, the District is looking at alternative locations near the school to begin air monitoring operations, while continuing to work with FUSD on allowing the air monitor to be installed, or should an agreement with the school district not be reached.

*Edison High School (Compact Multi-Pollutant System)*

The District has been in discussions with FUSD to place an Air Pointer compact multi-pollutant air monitoring system at Edison High School on the corner of California and Walnut Avenues. The District is continuing to attempt to schedule further meetings with FUSD to discuss implementation details. In the interim, the air monitoring van is being utilized to monitor areas nearby the school. In addition, the District is looking at alternative locations near the school to begin air monitoring operations, while continuing to work with FUSD on allowing the air monitor to be installed, or should an agreement with the school district not be reached.

*Yosemite Middle School (PM2.5)*

The District has been in discussions with FUSD in regards to placing a real-time PM2.5 monitor at Yosemite Middle School on the corner of 9<sup>th</sup> Street and Floradora Avenue. The District is continuing to attempt to schedule further meetings with FUSD to discuss implementation details. In the interim, the air monitoring van is being utilized to monitor areas nearby the school. In addition, the District is looking at alternative locations near the school to begin air monitoring operations, while continuing to work with FUSD on allowing the air monitor to be installed, or should an agreement with the school district not be reached.

*Malaga Elementary School (Air Monitoring Trailer)*

The District has been working with Fowler Unified School District to install the air monitoring trailer at Malaga Elementary School on the corner of Ward and Central Avenues. On February 26, 2020, the Fowler Unified School Board approved the proposed installation. District staff is now working with the school and electrical contractors to install the necessary electrical power for the air monitoring trailer. In the interim, the air monitoring van is being utilized to monitor areas nearby the school.

*Madison Elementary School (PM2.5)*

The District has been in discussions with Central Unified School District to place a real-time PM2.5 monitor at the school on the corner of S. Brawley Avenue and W. Madison Avenue. The District is still awaiting approval from Central Unified School District. In the interim, the air monitoring van is being utilized to monitor areas nearby this location. In addition, the District is looking at alternative locations near the school to begin air monitoring operations while details continue to be developed with Central Unified School District, or should an agreement with the school district not be reached.



### Mobile Air Monitoring Van Routes

In addition to the semi-mobile and fixed platforms, the District has been maximizing the usage of the considerable air monitoring capabilities of the air monitoring van to measure a variety of air pollutants of concern throughout the community. Measurements taken with the air monitoring van will allow the District and the community steering committee to understand local air pollution in the communities while also giving the District the ability to rapidly respond to air pollution concerns in other unmonitored regions. Intensive air monitoring operations with the mobile van began in January 2020. As mentioned earlier, the air monitoring van has enabled the District to commence air monitoring activities in areas that are still awaiting approval for installation of semi-mobile and fixed air monitoring equipment.

## II. Summary of PM<sub>2.5</sub> Air Monitoring

During this reporting period, concentrations of hourly PM<sub>2.5</sub> were measured at the air monitoring sites of Roosevelt High School, Bitwise South Stadium, and Fresno-Foundry within the South Central Fresno community boundary. The results of these measurements are summarized in the following table, and displayed in more detail in Appendix A to this report. Overall, through the period these sites experienced very few days when the federal 24-hour average standard of 35 µg/m<sup>3</sup> was exceeded, with most of these exceedances being influenced by fireworks emissions. PM<sub>2.5</sub> measurements across the community were fairly consistent through the period.


### **Quarterly 24-Hour Average PM<sub>2.5</sub> (µg/m<sup>3</sup>)**

<b>Quarter</b>	<b>Roosevelt High School</b>	<b>Bitwise South Stadium</b>	<b>Fresno-Foundry*</b>
2019 Q2	6.6	-	-
2019 Q3	5.7	6.2	-
2019 Q4	14.2	14.3	-
2020 Q1	13.7	12.5	14.7

\* Microscale site located adjacent to Highway 99


The following provides a comparison of daily PM<sub>2.5</sub> measurements during this period between the sites of Roosevelt High School, Bitwise South Stadium, Fresno-Foundry, and the nearby Clovis and Fresno-Garland air monitoring sites just north of the community boundary.

### 2019 Q2 Daily PM2.5 at Roosevelt High School, Clovis, and Fresno-Garland




Site	Average PM2.5 Concentration (µg/m <sup>3</sup> )
Roosevelt High School	6.6
Clovis	6.6
Fresno-Garland	7.5

### 2019 Q3 Daily PM2.5 at Roosevelt High School, Bitwise South Stadium, Clovis, and Fresno-Garland




Site	Average PM2.5 Concentration (µg/m <sup>3</sup> )
Roosevelt High School	5.7
Bitwise South Stadium	6.2
Clovis	8.9
Fresno-Garland	7.9

**2019 Q4 Daily PM2.5 at Roosevelt High School, Bitwise South Stadium, Clovis, and Fresno-Garland**



Site	Average PM2.5 Concentration ( $\mu\text{g}/\text{m}^3$ )
Roosevelt High School	14.2
Bitwise South Stadium	14.3
Clovis	11.6
Fresno-Garland	15.6

**2020 Q1 Daily PM2.5 at Roosevelt High School, Bitwise South Stadium, Fresno-Foundry, Clovis, and Fresno-Garland**



Site	Average PM2.5 Concentration ( $\mu\text{g}/\text{m}^3$ )
Roosevelt High School	13.7
Bitwise South Stadium	12.5
Fresno-Foundry*	14.7
Clovis	10.8
Fresno-Garland	13.1

\* Microscale site located adjacent to Highway 99

### III. Summary of Data Collected using Mobile Air Monitoring Van

In addition to the expanded PM2.5 measurements that have been occurring in the South Central Fresno community, the District has been taking advantage of the considerable air monitoring capabilities of the mobile air monitoring van to measure a variety of air pollutants throughout the community

The air monitoring van is ideal for taking measurements in unmonitored areas of concern and for regular surveillance over the entire community of South Central Fresno in short timeframes. The air monitoring van has also enabled the District to commence air monitoring activities in areas that are still awaiting approval for installation of semi-mobile and fixed air monitoring equipment.

In January 2020, the air monitoring van was used to measure a variety of air pollutants within the South Central Fresno community in the map below. The blue boundary in the map was identified by the South Central Fresno community steering committee as an area of interest for taking air quality measurements with the mobile air monitoring van.

### **CSC Recommended Areas of Interest for using Mobile Air Monitoring Van**

**Site A:** Parking lot on east corner of Tulare and 'R' Street


**Site B:** Parking lot on south east corner of East Butler Ave and South Cedar Ave

**Site C:** Parking lot on southwest corner of East California Ave and South Van Ness Ave

**Site D:** On 2<sup>nd</sup> Street south of intersection with Jensen Ave

**Site E:** Parking lot on southeast corner of E Jensen Ave and S Cedar Ave

**Site F:** Unpaved lot on southwest corner of E North Ave and S Cherry Ave



Following the January 2020 air monitoring van campaign described above, in March 2020 the District focused its efforts using the air monitoring van to monitor emissions near the aforementioned school sites that are awaiting approval for installation of semi-mobile and fixed air monitoring equipment.

The concentrations of pollutants measured with the air monitoring van at all locations described above were well below federal air quality standards during these periods, as summarized below. Note that although BTEX compounds were being measured by the air monitoring van during this period, concentrations of these compounds were not high enough to be detected by the analyzer.

### **Average and Peak Pollutant Concentrations in Areas Monitored with the Mobile Air Monitoring Van (January 2020, March 2020)**

Pollutant	Overall Average Value	Peak 1-hr Average Value	Applicable Standard
BTEX	None detected	None detected	-
PM <sub>2.5</sub>	4.8 µg/m <sup>3</sup>	16.6 µg/m <sup>3</sup>	35 µg/m <sup>3</sup> (24-hr average)
Ozone	24 ppb	53 ppb	70 ppb (8-hr average)
CO	0.25 ppm	0.59 ppm	35 ppm (1-hr average)

NO2	12 ppb	52 ppb	100 ppb (1-hr average)
SO2	0.5 ppb	0.9 ppb	75 ppb (1-hr average)

Appendix B to this report includes more details of daily measurements at each location using the mobile air monitoring van during the January and March 2020 timeframes.

#### IV. Summary of PM2.5 Speciation Analysis

To build a fuller understanding of the various constituents that comprise the overall PM2.5 concentrations in the South Central Fresno community, and their relative comparison, in November 2019 the District began operating PM2.5 speciation sampling at the Fresno-Foundry site near the intersection of Jensen Avenue and Highway 99. The collected samples were sent to a third-party laboratory for analysis to determine the contribution of various species of PM2.5 to the overall measured PM2.5 mass.

The nature and formation of PM2.5 in the San Joaquin Valley is highly complex as it can be composed of any material that has a diameter of 2.5 microns or less. PM2.5 can be emitted directly as primary PM2.5 from various sources or formed secondarily through chemical reactions in the atmosphere. The resulting ambient PM2.5 mixture can include aerosols (fine airborne solid particles and liquid droplets) consisting of components of nitrates, sulfates, organic carbon, black carbon, soil, trace metals, and more.


PM2.5 in the Valley is comprised of many species that contribute to the total PM2.5 mass. This complex mixture is attributable to emissions from stationary, mobile, and area-wide sources, as well as naturally occurring emissions. Although the list of species contributing to PM2.5 in the Valley is lengthy, it can be grouped into larger representative categories. The following is a brief description of each of these larger species categories:

- **Ammonium Nitrate:** Ammonium nitrate is formed from the reaction of ammonia and nitric acid, where the nitric acid is formed from emissions of nitrogen oxides.
- **Ammonium Sulfate:** Ammonium sulfate is formed from the reaction of ammonia and sulfuric acid, where the sulfuric acid is formed primarily from emissions of sulfur oxide, with smaller amounts forming from direct emissions of sulfur.
- **Organic carbon:** Organic carbon (OC) are generated as primary organic aerosol, predominantly through the combustion of hydrocarbons. Key sources include cooking, industrial processes, mobile source exhaust, tire wear, and wood burning. Secondary organic aerosols are formed from the oxidation of motor vehicle hydrocarbons, wood burning, solvent use, and industrial processes.


- Black Carbon:** Black carbon is also known as soot or elemental carbon, and is formed during incomplete combustion in fuels, including mobile exhaust (mainly diesel) and wood burning.
- Soil:** This category consists of road dust and soil dust that are entrained in the air from activity, such as soil disturbance or airflow from traffic.
- Other Metals:** Identified as components from soil emissions or found in other particulates having been emitted in connection with combustion from engine wear, brake wear, and similar processes. Certain metals are also emitted from the use of fireworks.

The following figures show the concentration levels and relative comparison of the various PM2.5 species sampled at the Fresno-Foundry air monitoring site. Samples were taken over a 4 month period during the timeframe of this report, and with the exception of a single day, the 24-hour average PM2.5 concentrations met the 24-hour federal PM2.5 standard of 35 µg/m<sup>3</sup>.

***Speciated PM2.5 Concentrations at Fresno-Foundry Site (November 2019 – March 2020)***



**Relative Comparison of PM<sub>2.5</sub> Species Measured at Fresno-Foundry Site  
 (November 2019 – March 2020)**



**V. Summary of VOC Speciation Analysis**

To build a fuller understanding of the various compounds that contribute to VOC concentrations in the South Central Fresno community, in December 2019 the District began operating VOC speciation sampling at the Fresno-Foundry site near the intersection of Jensen Avenue and Highway 99. The collected samples were sent to a third party laboratory for analysis to determine the various specific VOCs that were detected in the atmosphere. This laboratory analysis is able to isolate the concentrations of nearly 70 different VOCs from each air sample collected in the field.


VOCs are carbon chained compounds that vaporize in ambient conditions. Among these compounds include but, are not limited to, BTEX, 1,3-butadiene, PAH, aldehydes, naphthalene, and diethanolamine. These compounds are typically emitted from products such as paints, inks, organic solvents, petroleum products as well as vehicle exhaust. The health effects of these compounds vary but, long term exposure can have lasting adverse health effects. A more detailed list of possible VOCs and their health effects is provided by the California Office of Environmental Health Hazard Assessment (OEHHA)<sup>1</sup>.

<sup>1</sup> <https://oehha.ca.gov/air/general-info/oehha-acute-8-hour-and-chronic-reference-exposure-level-rel-summary>




During this period, the District collected 22 air samples for laboratory analysis. The VOC laboratory analysis is capable of isolating concentrations of 68 VOC species, however during this period most VOCs were not detected in the atmosphere. The following chart displays the small number of VOCs that were detected during this period, with most of these reporting only trace levels.

### ***VOC Species Detected at Fresno-Foundry Site (December 2019 – March 2020)***



During this period, methanol, ethanol, and acetone were the primary VOCs detected. Of these three, only methanol has an associated Reference Exposure Level (REL), a health risk metric established by the OEHHA. A comparison between the measured methanol concentrations and the associated REL is given in the following figure. Methanol is released to the environment during industrial uses and naturally from volcanic gases, vegetation, and microbes. It is released into ambient air from its evaporation during solvent uses or from automobile exhaust. The concentrations of methanol detected were well below the OEHHA REL chronic value, which is a more protective value than the acute REL. Overall, during the monitoring period no concerning concentrations of VOCs were detected in the samples taken.

## 24-Hr Methanol Concentrations



## VI. Availability and Access to Community Air Monitoring Data


Specific air quality data collected within the South Central Fresno community air monitoring network is available in real-time at the community air monitoring page located at <http://community.valleyair.org/selected-communities/south-central-fresno/air-monitoring/>. This District will also be posting to this page reports summarizing the laboratory speciation analysis conducted for PM<sub>2.5</sub> and VOCs. This District will continue to develop and publish quarterly reports summarizing collected data to the District's AB 617 air monitoring website. Collected community air monitoring data is also available for download on the California Air Resources Board (CARB) AQview tool located at <https://ww2.arb.ca.gov/es/community-air-quality-portal>, where collected air monitoring data from all AB 617 communities is uploaded.

## VII. Next Steps


The District has deployed air monitoring equipment and obtained real-time PM<sub>2.5</sub> measurements along with numerous sample collections for advanced third-party laboratory speciation analyses of PM<sub>2.5</sub> and VOC for a large range of pollutants throughout the South Central Fresno community. Emissions concentrations measured throughout this period were predominantly below federal standards and OEHHA REL values. Moving forward, the District will continue these measurements and will continue to expand and establish the community air monitoring network in the community of South Central Fresno.

### VIII. Appendix A: Daily PM2.5 Averages


#### Roosevelt High School – March 2019




#### Roosevelt High School – April 2019




***Roosevelt High School – May 2019***




***Roosevelt High School – June 2019***




**Roosevelt High School – July 2019**




**Roosevelt High School – August 2019**




***Roosevelt High School – September 2019***




***Roosevelt High School – October 2019***




Roosevelt High School – November 2019




Roosevelt High School – December 2019



**Roosevelt High School – January 2020**




**Roosevelt High School – February 2020**







**Roosevelt High School – March 2020**




**Bitwise South Stadium – August 2019**




**Bitwise South Stadium – September 2019**




**Bitwise South Stadium – October 2019**




***Bitwise South Stadium – November 2019***




***Bitwise South Stadium – December 2019***




***Bitwise South Stadium – January 2020***



***Bitwise South Stadium – February 2020***



***Bitwise South Stadium – March 2020***



## IX. Appendix B: Summary of Data Collected using Mobile Air Monitoring Van

During the period of January 1 to January 22, 2020, the mobile air monitoring van was used to measure a variety of pollutants at the following locations within the South Central Fresno community.

**Site A:** Parking lot on east corner of Tulare and 'R' Street


**Site B:** Parking lot on south east corner of East Butler Ave and South Cedar Ave

**Site C:** Parking lot on southwest corner of East California Ave and South Van Ness Ave

**Site D:** On 2<sup>nd</sup> Street south of intersection with Jensen Ave

**Site E:** Parking lot on southeast corner of E Jensen Ave and S Cedar Ave

**Site F:** Unpaved lot on southwest corner of E North Ave and S Cherry Ave



During the period of March 6 to March 26, 2020, the air monitoring van coverage was extended to areas that are still awaiting approval for installation of semi-mobile and fixed air monitoring equipment. During the days during this period, the air monitoring van was kept in a stationary mode for multiple hours to capture a longer period of pollutant trends in the area. These areas include locations near Orange Center Elementary School, Heaton Elementary School, Yosemite Middle School, Edison High School, and Madison Elementary School.

Note that although BTEX compounds were being measured by the air monitoring van during this period, concentrations of these compounds were not high enough to be detected by the analyzer. The measured pollutant concentrations are detailed in the following tables. Note that CO concentrations are reported in parts per million, PM<sub>2.5</sub> is reported in micrograms per cubic meter, while all other pollutants are reported in parts per billion.

January 7, 2020

Site	Time	O3	CO	NO2	SO2	H2S
A	09:37-10:15	7.1	0.59	20.8	0.42	0
B	10:26-11:05	10.9	0.58	13.3	0.7	0
C	11:13-11:55	9.68	0.49	23.5	0.9	0
D	11:58-12:49	9.56	0.52	28.3	0.6	0
E	12:55-13:30	14.2	0.53	19.5	0.5	0
F	13:35-14:20	10.1	0.3	52.1	0.5	0

January 15, 2020

Site	Time	O3	CO	NO2	SO2	H2S
A	12:45-13:23	27	0.29	12.9	1.71	0.2
B	11:58-12:38	27.1	0.39	6.43	1.6	0.5
C	11:14-11:50	17.3	0.37	23.2	1.9	1.0
D	10:32-11:06	14.4	0.7	24.2	1.9	1.5
E	09:41-10:27	9.0	0.83	37.5	2.8	2.6
F	08:56-09:33	2.7	0.6	34.7	2.6	3.3


January 22, 2020

Site	Time	O3	CO	NO2	SO2	H2S
A	09:13-09:47	10	0.13	16.9	0.79	1.9
B	09:53-10:25	11.2	0.12	15.5	12.0	1.8
C	10:33-11:04	16.0	0.03	21.0	0.1	1.7
D	11:13-11:51	25.1	0.02	12.2	0.0	1.4
E	11:55-12:32	43.3	0.32	19.3	1.8	2.3
F	10:16-10:52	25.6	0.04	12.4	0.3	1.3


March 6, 2020 Site B

Time	O3	CO	NO2	NO	NOx	H2S	SO2	PM2.5
9:00:00	18.9	0.42	21.3	16.6	38.0	0.0014	0.0003	7.5
10:00:00	24.2	0.42	25.8	18.6	44.3	0.0015	0.0005	23.0
11:00:00	40.8	0.27	9.1	3.9	13.0	0.0010	0.0006	16.2
12:00:00	46.6	0.23	8.5	3.0	11.5	0.0006	0.0005	13.4
13:00:00	53.2	0.19	6.4	2.1	8.5	0.0005	0.0005	10.6
14:00:00	50.4	0.31	10.1	4.3	14.3	0.0004	0.0005	5.5

### PM2.5



### Ozone



### NO2







March 2020 Near Orange Center Elementary School

Date	Time	O3	CO	NO2	NO	NOx	H2S	SO2	PM2.5
03/10/2020	8:00	5.3	0.24	19.0	15.3	34.3	0.0022	0.0000	3.3
	9:00	4.9	0.24	20.9	12.4	33.3	0.0021	0.0005	3.6
	10:00	6.9	0.22	19.2	12.7	31.9	0.0019	0.0007	6.5
	11:00	8.8	0.20	10.8	4.5	15.3	0.0010	0.0002	3.9
	12:00	9.5	0.20	9.9	3.3	13.2	0.0010	0.0002	3.9
	13:00	10.9	0.22	12.9	5.0	17.9	0.0007	0.0002	3.5
	14:00	12.4	0.27	12.6	7.3	19.9	0.0003	0.0002	2.4
03/12/2020	7:00	-	0.80	26.4	71.2	97.7	0.0014	0.0008	4.7
	8:00	-	0.31	13.0	20.1	33.2	0.0027	0.0006	14.4
	9:00	-	0.23	8.3	8.6	16.9	0.0027	0.0007	16.7
	10:00	-	0.21	4.8	3.5	8.4	0.0017	0.0006	16.8
	11:00	-	0.20	5.9	4.4	10.4	0.0013	0.0005	15.4
	12:00	-	0.19	8.7	4.1	12.7	0.0009	0.0006	15.6
03/19/2020	9:00	-	0.18	7.5	3.8	11.4	0.0004	0.0001	4.0
	10:00	-	0.17	1.9	2.9	4.8	0.0005	0.0002	2.7
	11:00	-	0.14	2.7	3.7	6.4	0.0003	0.0002	3.2
	12:00	-	0.14	12.5	6.6	19.1	0.0005	0.0003	4.4
	13:00	-	0.12	1.7	1.6	3.4	0.0004	0.0003	3.7
	14:00	-	0.16	2.9	4.0	6.9	0.0002	0.0001	3.4
	15:00	-	0.21	6.7	6.1	12.8	0.0003	0.0002	3.1

PM2.5




Ozone



Community of South Central Fresno


NO2




*March 16, 2020 Near Heaton Elementary School*

Time	O3	CO	NO2	NO	NOx	H2S	SO2	PM2.5
9:00	-	0.16	3.3	1.5	4.8	0.0005	0.0000	1.4
10:00	-	0.15	3.1	1.2	4.2	0.0011	0.0002	1.7
11:00	-	0.17	2.5	1.1	3.7	0.0007	0.0001	1.4
12:00	-	0.17	5.4	16.5	22.0	0.0005	0.0002	0.8
13:00	-	0.16	17.0	1.1	18.1	0.0005	0.0001	1.3
14:00	-	0.17	16.2	1.0	17.2	0.0003	0.0001	1.0

PM2.5




NO2




March 16, 2020 Near Yosemite Middle School

Time	O3	CO	NO2	NO	NOx	H2S	SO2	PM2.5
8:00	-	0.19	3.0	3.2	6.2	0.0003	0.0001	1.5
9:00	-	0.16	3.3	1.5	4.8	0.0005	0.0000	1.4
10:00	-	0.15	3.1	1.2	4.2	0.0011	0.0002	1.7
11:00	-	0.17	2.5	1.1	3.7	0.0007	0.0001	1.4
12:00	-	0.17	5.4	16.5	22.0	0.0005	0.0002	0.8
13:00	-	0.16	17.0	1.1	18.1	0.0005	0.0001	1.3
14:00	-	0.17	16.2	1.0	17.2	0.0003	0.0001	1.0
15:00	-	0.22	8.6	3.6	12.2	0.0004	0.0003	0.6

PM2.5




NO2




March 19, 2020 Near Edison High School

Time	O3	CO	NO2	NO	NOx	H2S	SO2	PM2.5
9:00	32.9	0.09	5.9	2.6	8.5	0	0	0.7
10:00	36.4	0.12	3.4	1.4	4.8	0	0	0.6
11:00	38.7	0.19	4.0	1.1	5.1	0	0	0.5
12:00	38.1	0.17	3.6	0.9	4.5	0	0	0.7
13:00	40.2	0.15	3.1	0.8	3.9	0	0	0.7


PM2.5



NO2




Ozone




March 26, 2020 Near Madison Elementary School

Time	O3	CO	NO2	NO	NOx	H2S	SO2	PM2.5
8:00:00	30.2	0.19	3.3	1.5	4.7	0	0.003	3
9:00:00	35.1	0.19	3.0	1.5	4.5	0	0.005	5
10:00:00	40.0	0.15	1.8	1.2	3.0	0	0.005	5
11:00:00	41.4	0.15	1.9	1.3	3.2	0	0.001	1
12:00:00	43.0	0.17	2.0	1.1	3.1	0	0	0
13:00:00	43.5	0.20	1.5	0.7	2.3	0	0	0

PM2.5



NO2



Ozone

